




Manuela Nagel (Autor)
Maintenance of plant genetic resources
Towards an understanding of seed deterioration



<https://cuvillier.de/de/shop/publications/8602>

Copyright:
Cuvillier Verlag, Inhaberin Annette Jentsch-Cuvillier, Nonnenstieg 8, 37075 Göttingen,
Germany
Telefon: +49 (0)551 54724-0, E-Mail: info@cuvillier.de, Website: <https://cuvillier.de>

INDEX

| | |
|---|-----|
| INDEX OF ABBREVIATIONS | III |
| 1 INTRODUCTION | 1 |
| 2 PROLOG | 2 |
| 2.1 Maintenance of plant genetic resources in <i>ex situ</i> gene banks | 2 |
| 2.1.1 The importance to collect plant genetic resources | 2 |
| 2.1.2 Long-term storage and utilization of <i>ex situ</i> collections | 3 |
| 2.1.3 Wheat genetic resources | 6 |
| 2.1.4 Barley genetic resources | 7 |
| 2.1.5 Oil seed rape genetic resources | 8 |
| 2.2 Survival in the dry state | 9 |
| 2.3 The establishment of seed longevity during seed maturation | 11 |
| 2.4 The physics of water and water activity in seeds | 15 |
| 2.5 The glass formation as requirement for seed desiccation and storability | 19 |
| 2.6 Seed viability loss during storage | 21 |
| 2.6.1 Prediction and factors of seed viability loss | 21 |
| 2.6.2 Biochemical mechanisms of seed viability loss | 24 |
| 2.7 Seed protection mechanisms during quiescence | 26 |
| 2.8 Aim of the study | 31 |
| 3 SCIENTIFIC PUBLICATIONS | 32 |
| 3.1 PAPER 1: The longevity of crop seeds stored under ambient conditions | 32 |
| 3.2 PAPER 2: Machine learning links seed composition, glucosinolates and viability of oilseed rape after 31 years of long-term storage | 45 |
| 3.3 PAPER 3: Changes in tocochromanols and glutathione reveal differences in the mechanisms of seed ageing under seedbank conditions and controlled deterioration in barley | 55 |
| 3.4 PAPER 4: Wheat seed ageing viewed through the cellular redox environment and changes in pH | 64 |
| 3.5 PAPER 5: Comparative physiology and proteomics of two wheat genotypes differing in seed storage tolerance | 92 |

| | | |
|------|---|-----|
| 3.6 | PAPER 6: Ageing is highly associated with lipid oxidation and hydrolysis in wheat seeds | 102 |
| 3.7 | PAPER 7: Barley seed ageing: genetics behind the dry elevated pressure of oxygen ageing and moist controlled deterioration. | 124 |
| 3.8 | PAPER 8: Seed longevity in oilseed rape (<i>Brassica napus</i> L.) - genetic variation and QTL mapping | 136 |
| 3.9 | PAPER 9: Genetic variation for secondary seed dormancy and seed longevity in a set of black-seeded European winter oilseed rape cultivars | 141 |
| 3.10 | PAPER 10: QTL analysis of falling number and seed longevity in wheat (<i>Triticum aestivum</i> L.) | 148 |
| 3.11 | PAPER 11: Effects of <i>Rht</i> dwarfing alleles on wheat seed vigour after controlled deterioration | 157 |
| 3.12 | PAPER 12: Genetic architecture of seed longevity in bread wheat (<i>Triticum aestivum</i> L.) | 166 |
| 3.13 | PAPER 13: Novel loci and a role for nitric oxide for seed dormancy and pre-harvest sprouting in barley | 176 |
| 4 | EPILOG | 165 |
| 4.1 | Seed deterioration processes are determined by the storage conditions | 165 |
| 4.2 | Differences in seed longevity are the consequence of compositional variations | 170 |
| 4.3 | Maternal growth environment influences seed longevity and seed dormancy | 174 |
| 4.4 | Molecular markers for seed deterioration at different water contents | 176 |
| 4.5 | The implications for gene bank management | 180 |
| 5 | SUMMARY AND CONCLUSIONS | 182 |
| 6 | REFERENCES OF PROLOG AND EPILOG | 184 |
| | ACKNOWLEDGEMENTS | 196 |
| | CURRICULUM VITAE | 197 |